arXiv:1406.3752v3 [math.AG] 28 Jul 2015

# Rationally connected non-Fano type varieties

Igor Krylov

**Abstract**

Varieties of Fano type are very well behaved with respect to MMP, and they are known to be rationally connected. We study the relation between classes of rationally connected varieties and varieties of Fano type. It is known that these classes are birationally equivalent in dimension 2. We give examples of rationally connected varieties of dimension $\geqslant 3$ which are not birational to Fano type varieties, thereby answering the question of Cascini and Gongyo [2, Question 5.2].

## 1 Introduction

The log minimal model program (MMP) is one of the key notions in birational geometry. Finding when can we run the MMP for a pair $(X, D)$ ($D$-MMP) is one of the central subjects and is still being developed. If $X$ is a variety of Fano type then we can run the $D$-MMP on it for any effective divisor $D$ on $X$ [15, Corollary 2.7]. We say that a normal projective variety $X$ is of Fano type if there is an effective $\mathbb{Q}$-divisor $H$ on $X$ such that the pair $(X, H)$ is Kawamata log terminal and $-(K_X + H)$ is ample. Varieties of Fano type have been introduced by Shokurov and Prokhorov in [15]. The Fano type property is preserved under flips and contractions ([15, Lemma 2.8]) therefore at any stage of the minimal model program on a variety of Fano type we have a Fano type variety. In particular the result of running the MMP on a variety of Fano type is a variety of Fano type. Varieties of Fano type appear naturally as quotients of Fano varieties and as resolutions of singular Fano varieties.

It would be nice if we could say if in a given birational class there is a variety which behaves well under $D$-MMP for any effective divisor $D$. For example a variety of Fano type. On the other hand it is known that varieties of Fano type are rationally connected [23, Theorem 1]. Therefore it is natural to ask if the converse is true. It is not even in dimension 2, consider a blow up of 10 points in general position on $\mathbb{P}^2$, it is not a variety of Fano type. The following question however makes sense.

**Question 1** ([2, Question 5.2])**.** *Let $X$ be a rationally connected variety. Is $X$ birationally equivalent to a variety of Fano type?*

In dimension 2 the answer is positive since every rationally connected surface is rational. In this paper we construct examples for which the answer is negative in dimension $\geqslant 3$. Namely the purpose is to prove the following theorem.

**Theorem 1.1.** *(i) Let $W$ be a generic smooth divisor of degree $(2M, 2l)$ on $\mathbb{P}^M \times \mathbb{P}^1$, $M \geqslant 3$, $l \geqslant 3$. Let $V$ be a double cover ramified over $W$. Let $\pi : V \dashrightarrow Y$ be a birational map to a Mori fiber space $f : Y \to Z$. Then $Z = \mathbb{P}^1$, $\pi$ is an isomorphism and the following diagram is commutative*

$$\begin{array}{ccc} V & \xrightarrow{\pi} & Y \\ \downarrow & & \downarrow f \\ \mathbb{P}^1 & \xrightarrow{g} & Z \end{array}$$

*(ii) Let $Y = \big(\mathbb{C}^6 \setminus Z\big(\langle u, v\rangle \cap \langle x, y, z, w\rangle\big)\big)/(\mathbb{C}^*)^2$, where $(\mathbb{C}^*)^2$-action is given by a matrix*

$$\begin{pmatrix} u & v & x & y & z & w \\ 0 & 0 & 1 & 1 & 2 & 3 \\ 1 & 1 & -3 & -3 & 0 & 0 \end{pmatrix}$$

*and let*

$$Q = w^2 + z^3 + (u^{12} + v^{12})M_4(x, y)z + R_{18}(u, v)x^2y^2(x - y)^2,$$

*where $M_4 \in \mathbb{C}[x, y]$, $\deg M_4 = 4$ and $R_{18} \in \mathbb{C}[u, v]$, $\deg R_{18} = 18$. Let $\mathcal{L}$ be a linear system on $Y$ given by $Q = 0$. Let $X$ be a generic element of $\mathcal{L}$. Let $\pi : X \dashrightarrow Y$ be a birational map to a Mori fiber space $f : Y \to Z$. Then $Z = \mathbb{P}^1$, $\pi$ is an isomorphism and the following diagram is commutative*

$$\begin{array}{ccc} X & \xrightarrow{\pi} & Y \\ \downarrow & & \downarrow f \\ \mathbb{P}^1 & \xrightarrow{g} & Z \end{array}$$

The varieties $V$ are Fano fibrations over $\mathbb{P}^1$. We say that a Fano fibration $U$ over $\mathbb{P}^1$ is birationally superrigid (Definition 2.4) if any birational

map $f : U \to Y$ to a Mori fiber space $Y \to Z$ is a fiberwise map, that is the following diagram is commutative

$$\begin{array}{ccc} U & \overset{\pi}{\dashrightarrow} & Y \\ \downarrow & & \downarrow f \\ \mathbb{P}^1 & \overset{g}{=\!=} & Z \end{array}$$

In particular it means that $U$ is not birational to a $\mathbb{Q}$-factorial terminal Fano variety or a conic bundle and is not rational. First, we use the result of Pukhlikov ([19, Theorem 2.1]) to show that $V$ is birationally superrigid. Then we use Tian's $\alpha$-invariant to conclude that any map to a Mori fiber space is an isomorphism.

Varieties $X$ are del Pezzo fibrations of degree 1. A priory we may have non-trivial maps to other del Pezzo fibrations of degree 1, fiberwise transformations [10], therefore we have to use different approach. By Lefschetz-type result [20] that $X$ is a Mori fiber space which allows us to use techinques of birational rigidity ([16], [6]). Suppose $f : X \to X'$ is a birational map which is not an isomorphism and $X'$ is a Mori fiber space. It follows from rigidity techniques that there is a linear system $\mathcal{M}$ with bad singularities on $X$. We note that a locus of bad singularities of $\mathcal{M}$ must contain a cusp $P$ of a rational curve on the fiber of anticanonical degree 1. Then we prove that all such points are the ordinary double points of $X$ or of the fibers on $X$. We show that a pair $(X, \mathcal{M})$ is always good at ordinary double points of $X$ or of fibers of $X$. At last we conclude that all birational maps from $X$ to Mori fiber spaces are isomorphisms.

We prove that any variety of Fano type is birationally equivalent to a Mori fiber space with big anticanonical divisor (Proposition 5.1). As $-K_V$ and $-K_X$ are not big (Lemmas 3.8 and 4.6) we get the following statement:

**Corollary 1.2.** *Varieties $V$ and $X$, described in Theorem 1.1, are not birationally equivalent to a variety of Fano type.*

Since $V$ and $X$ are rationally connected (Corollary 5.4) we conclude that the answer to Question 1 is negative for these varieties.

## Acknowledgements

The author would like to thank John Ottem and Ivan Cheltsov, for bringing this question to his attention and for many helpful conversations, and Antony Macioca, Milena Hering, Johan Martens, and Michael Wemyss for useful suggestions. The author was supported by an Edinburgh PCDS scholarship.

# 2 Preliminaries

All varieties in this paper are considered to be normal, projective $\mathbb{Q}$-factorial and defined over $\mathbb{C}$ unless stated otherwise. By a Fano fibration over a projective line we mean a terminal $\mathbb{Q}$-factorial variety with Picard number 2 and a map to a projective line such, that the generic fiber is a smooth Fano variety. For a definition of Mori fiber space in general we refer to [1]. We write $\sim_{\mathbb{Q}}$ for linear equivalence of $\mathbb{Q}$-divisors.

**Definition 2.1** ([14, p. 6])**.** Let $X$ be a normal variety and $D$ be a $\mathbb{Q}$-divisor on $X$ such that $K_X + D$ is $\mathbb{Q}$-Cartier. Let $\pi : \widetilde{X} \to X$ be a birational morphism from a normal variety $\widetilde{X}$ and let $\widetilde{D} = \pi^{-1}(D)$ be a proper transform of $D$. Then we can write

$$K_{\widetilde{X}} + \widetilde{D} \sim_{\mathbb{Q}} \pi^*(K_X + D) + \sum_E a(E, X, D)E,$$

where $E$ runs through all the distinct exceptional divisors of $\pi$ on $\widetilde{X}$ and $a(E, X, D)$ is a rational number. We say that the pair $(X, D)$ is *terminal* (resp. *canonical*, *log terminal*, *log canonical*) if $a(E, X, D) > 0$ (resp. $a(E, X, D) \geqslant 0$, $a(E, X, D) > -1$, $a(E, X, D) \geqslant -1$) for every prime $\pi$-exceptional divisor $E$ on any normal variety $\widetilde{X}$ such that there is a birational morphism from $\widetilde{X}$ to $X$.

If $D = 0$, we simply say that $X$ has only terminal (resp. canonical, log terminal, log canonical) singularities. The numbers $a(E, X) = a(E, X, 0)$ are called *discrepancies* of the divisor $E$. We say that the pair $(X, D)$ is *klt* if it is log terminal and coefficients $a_i$ of all prime divisors $D_i$ in $D = \sum a_i D_i$ are less than 1. Sometimes we will say that $K_X + D$ is terminal (resp. canonical, etc) instead of $(X, D)$ is terminal (resp. canonical, etc).

We say that the pair $(X, D)$ has terminal (resp. canonical, log terminal, log canonical) singularities at a subvariety $Z \subset X$ if for every birational morphism $f : X' \to X$ and for every exceptional divisor $E$ such that $f(E) = Z$ the inequality $a(E, X, D) > 0$ (resp. $a(E, X, D) \geqslant 0$, $a(E, X, D) > -1$, $a(E, X, D) \geqslant -1$) holds.

Let $\mathcal{M}$ be a linear system on $X$ and let $\lambda \in \mathbb{Q}$ be a non-negative number. We say that the pair $(X, \lambda\mathcal{M})$ has terminal (resp. canonical, klt, log canonical) singularities at $Z \subset X$ if for a generic divisor $D \in \mathcal{M}$ the pair $(X, \lambda D)$ is terminal (resp. canonical, klt, log canonical) at $Z$.

**Example 2.2.**
- Consider the pair $(S, C)$, where $S$ is a smooth surface and $C \subset S$ is a smooth curve. Then the pair is canonical at every point of $C$ and is terminal at every point other point.

- Consider the pair $(\mathbb{P}^2, |L|)$, where $L$ is a line. Then the pair is terminal at every point $P \in \mathbb{P}^2$ since a generic line does not pass through $P$.
- Let $\mathcal{L}$ be the linear system of lines on $\mathbb{P}^2$ passing through a point $P$. Then the pair $(\mathbb{P}^2, \mathcal{L})$ is canonical at $P$ and is terminal elsewhere.

**Definition 2.3** ([15, Lemma-Definition 2.6])**.** We say that a variety $X$ is of *Fano type* if there exists an effective $\mathbb{Q}$-divisor $D$ on $X$ such that $-(K_X + D)$ is $\mathbb{Q}$-Cartier and ample, and the pair $(X, D)$ is klt.

**Example 2.4.** (i) The quotient $Y = X/G$ of a Fano variety by a finite group is a variety of Fano type. Since $K_X = f^*(K_Y + \frac{R}{|G|})$, where $R$ is the ramification divisor, the divisor $-(K_Y + \frac{R}{|G|})$ is ample. The pair $(Y, \frac{R}{|G|})$ is klt [13, Proposition 3.16] hence $Y$ is a variety of Fano type.

(ii) Suppose $(S, D_S)$ is a klt del Pezzo surface. Naturally it is of Fano type. Let $f : X \to S$ be a minimal resolution of singularities of $S$ and let $D$ be a proper transform of $D_S$. Then $X$ is also of Fano type. Indeed we can write

$$K_X + D = f^*(K_S + D_S) + \sum a_i E_i,$$

where $-1 < a_i \leqslant 0$, thus $F = -(K_X + D + \sum a_i E_i)$ is a pullback of an ample divisor. Note that $F$ is not ample because $F \cdot E_i = 0$. Consider the pair $(X, D + \sum(1-\varepsilon)a_i E_i)$, for $\varepsilon << 0$ the pair is klt and $-K_X + D + \sum(1-\varepsilon)a_i E_i$ is ample.

(iii) Suppose a surface $S$ is of Fano type. Then there is a divisor $D$ such that $-(K_S + D)$ is ample. Thus $-K_S$ is the sum of ample and effective divisor, therefore it is big. If $S$ is a blow up of 9 points on $\mathbb{P}^2$ in general position then $-K_S$ is not big, therefore such $S$ is not of Fano type. Note that if points are not in a general position then $-K_S$ might still be big [21].

Fano type varieties also appear as exceptional divisors of extremal contractions.

**Definition 2.5** ([17, Definition 1])**.** We say that a Fano variety $V$ with $\mathrm{Pic}(V) = \mathbb{Z}$ is *birationally superrigid* if $\mathrm{Bir}(V) = \mathrm{Aut}(V)$. We say that a Fano fibration $f : V \to \mathbb{P}^1$ is *birationally superrigid* if for any Mori fiber space $f' : V' \to S$ and any birational map $\pi : V \dashrightarrow V'$ the base $S$ is $\mathbb{P}^1$, there is a commutative diagram

$$\begin{array}{ccc} V & \overset{\pi}{\dashrightarrow} & V' \\ \downarrow{\scriptstyle f} & & \downarrow{\scriptstyle f'} \\ \mathbb{P}^1 & \overset{g}{=\!=} & \mathbb{P}^1, \end{array}$$

and $\pi$ is an isomorphism on a generic fiber of $f$.

It follows from the definition that, if $V$ is a birationally superrigid fibration over $\mathbb{P}^1$, then $\pi$ is a fiberwise map.

Consider a $\mathbb{P}^1$-bundle $f : X \to \mathbb{P}^1$. We can blow up a point on a fiber $F$ of $f$ and then contract the proper transform of $F$. This is an elementary transformation of $\mathbb{P}^1$ bundles and this is the simplest example of a fiberwise map. Let us look at a more general situation. Consider a Fano fibration $f : X \to \mathbb{P}^1$, let $Z \subset F$ be a subvariety on a fiber of $F$. Denote the blow up of $Z$ as $\psi_Z : X_Z \to X$. Suppose we can do some flops $\rho : X_Z \dashrightarrow \widetilde{X}_Z$ in such a way that there is a divisorial contraction $\psi_F : \widetilde{X}_Z \to \widetilde{X}$ of the proper transform of $F$. The birational map $\psi_F \circ \rho \circ \psi_Z^-1$ is an elementary fiberwise map. One can look at fiberwise map as a composition of elementary fiberwise maps.

**Definition 2.6** ([3, Definition 1.2])**.** Let $X$ be a $\mathbb{Q}$-factorial Fano variety with log terminal singularities. Define *Tian's $\alpha$-invariant* of $X$ by the number

$$\alpha(X) = \sup\{c : (X, cD) \text{ is log canonical} \\ \text{for every effective } \mathbb{Q}\text{-divisor } D,\ D \sim_{\mathbb{Q}} -K_X\}.$$

Tian has introduced the $\alpha$-invariant for Kähler manifolds in [22] in a different way but it coincides with the one defined above ([5, Theorem A.3]).

**Proposition 2.7** ([6, Proposition 2.7])**.** *Let $X$, $X'$ be varieties with $\mathbb{Q}$-factorial singularities, and let $f : X \dashrightarrow X'$ be a birational map which is an isomorphism in codimension one. Let $H'$ be an ample divisor on $X$. If $H = f^{-1}H'$ is ample then $\Phi$ is an isomorphism.*

*Proof taken from [6].* Let $Y$ be a resolution of $f$ and let $p : Y \to X$ and $q : Y \to X'$ be the corresponding morphisms. Let $E_i$ be the $p$-exceptional divisors. Since $f$ is an isomorphism in codimension one, $q$ has the same exceptional divisors. For suitable $a_i$ we have

$$\sum a_i E_i \sim_{\mathbb{Q}} -p^*H + q^*H'.$$

It follows from the negativity lemma [11, Lemma 2.19] that $a_i = 0$ and $p^*H = q^*H'$. Let $C \subset Y$ be a curve such that $q(C) \subset X'$ is a point. By the projection formula

$$0 = q(C) \cdot H' = C \cdot q^*H' = C \cdot p^*H = p_*(C) \cdot H$$

hence $p(C)$ is a point since $H$ is ample. Thus $f$ is an isomorphism. □

**Proposition 2.8** (Noether-Fano inequality, [6, Theorem 4.2])**.** *Let $V$ be a terminal Fano variety with* $\operatorname{Pic} V = \mathbb{Z}$*. Then $V$ is birationally superrigid if and only if there are no movable linear systems $\mathcal{M}$ on $V$ such that $\lambda\mathcal{M} + K_V \sim_{\mathbb{Q}} 0$ and the pair $(V, \lambda\mathcal{M})$ is not canonical for some $\lambda \in \mathbb{Q}$.*

*Proof.* Let $V$ be birationally superrigid and let $\mathcal{M}$ be a movable linear system on $V$. Suppose that $\lambda\mathcal{M} + K_V \sim_{\mathbb{Q}} 0$ and the pair $(V, \lambda\mathcal{M})$ is not canonical for some $\lambda \in \mathbb{Q}$. Consider a resolution of singularities $h : \widetilde{V} \to V$ of $\mathcal{M}$ and the generic divisor $\widetilde{D} \in h^{-1}(\mathcal{M})$. Without loss of generality we may assume that the pair $(V, \lambda\widetilde{D})$ is terminal. Let $E_i$ denote the exceptional divisors of $h$. Let the pair $(V', \lambda D')$ be a result of the $\lambda\widetilde{D}$-MMP. A divisor $K_{V'} + \lambda D'$ is not nef since

$$K_{\widetilde{V}} + \lambda\widetilde{D} \sim_{\mathbb{Q}} \sum a_i E_i,$$

therefore $V'$ is a Mori fiber space. Let $g : V \dashrightarrow V'$ be the composition of $h^{-1}$ and the $\lambda D$-MMP. It follows from the superrigidity of $V$ that $g$ is an isomorphism and $D' = g(D)$. The pair $(V', \lambda g(D))$ is terminal as it is a result of the MMP but $(V, \lambda D)$ is not canonical thus we have a contradiction.

Suppose $g : V \dashrightarrow X$ is a birational map to a Mori fiber space $X$. First, suppose $X$ is not a Fano variety. Then there is a Fano fibration $\pi : X \to S$, where $\dim S \geqslant 1$. Let $H_X$ be the pullback of a very ample divisor on $S$ and consider the linear system $\mathcal{M}_X = \big|H\big|$. Put $\mathcal{M} = g^{-1}\mathcal{M}_X$. There is a $\lambda \in \mathbb{Q}$ such that $\lambda\mathcal{M} + K_V \sim_{\mathbb{Q}} 0$. Let $W$ be a resolution of $g$ and let $\psi : W \to V$, $\varphi : W \to X$ be the corresponding morphisms.

We can write on $W$

$$\begin{aligned}\psi^*(K_V + \lambda\mathcal{M}) + \sum a_i E_i &\sim_{\mathbb{Q}} K_W + \lambda\mathcal{M}_W \\ &\sim_{\mathbb{Q}} \varphi^*(K_X + \lambda\mathcal{M}_X) + \sum b_j E_j. \qquad (1)\end{aligned}$$

Note that a generic fiber $f$ of $\pi$ does not pass through centers of the exceptional divisors of $\varphi$ and $f \cdot H_X = 0$. Intersecting $\varphi^* f$ with (1) we get

$$\sum a_i(\varphi^* f \cdot E_i) \sim_{\mathbb{Q}} \varphi^*(f \cdot K_X).$$

Since $-K_X$ is ample on $f$ because $X$ is a Mori fiber space. On the other hand $\varphi^* f$ does not lie on $E_i$ because $f$ is generic, thus $E_i|_f$ is effective for all $i$. Therefore there is some $i$ such that $a_i < 0$.

Thus Mori fiber space $X$ must be a Fano variety, in particular $\operatorname{Pic}(X) = \mathbb{Z}$. Take a very ample divisor $H \in \big|-nK_X\big|$ on $X$, $n > 1$. Define $\mathcal{M}, \mathcal{M}_X, \lambda, W, \varphi, \psi$ as in previous case, then we can write (1) on $W$ again.

The linear system $\mathcal{M}_X$ is free, therefore for all $j$ , $b_j > 0$. It is easy to see that

$$K_X + \lambda \mathcal{M}_X \sim_{\mathbb{Q}} (1 - \lambda n) K_X.$$

Suppose that $1 - \lambda n > 0$ then by intersecting (1) with the pullback of a generic divisor $D \in \mathcal{M}_{\mathcal{X}}$ we have

$$\sum a_i (\varphi^*(D) \cdot E_i) \sim_{\mathbb{Q}} \varphi^*((1 - \lambda n) D \cdot K_X).$$

On the right-hand side we have the pullback of the intersection of an ample and an anti-ample class. On the left-hand side we have an intersection of effective divisors with the pullback of an ample class, thus there is an $i$ such that $a_i < 0$.

Suppose that $1 - \lambda n \leqslant 0$ and $a_i \geqslant 0$ for all $i$. Rewriting (1) we have

$$\varphi^*((1 - \lambda n) K_X) \sim_{\mathbb{Q}} \sum a_i E_i - \sum b_j E_j.$$

Suppose there $i$ such that $E_i$ is not an exceptional divisor of $\varphi$. Then by negativity lemma $-b_i \geqslant 0$. Thus, all exceptional divisors of $\psi$ are also exceptional for $\varphi$. Since the Picard numbers of $V$ and $X$ are equal to 1, morphisms $\psi$ and $\varphi$ have the same number of exceptional divisors. Therefore $\psi$ and $\varphi$ have the same exceptional divisors hence $g$ is an isomorphism in codimension 1. By Proposition 2.7, $g$ is an isomorphism. □

**Remark 2.9.** To prove that some variety is superrigid we need to show that $(X, D)$ is a canonical pair for any $D \sim_{\mathbb{Q}} -K_X$. Thus method works best for varieties with "small" $-K$: hypersurfaces of degree $n$ in $\mathbb{P}^n$ and double covers of $\mathbb{P}^n$ ramified over hypersurface of degree $2n$. In high dimension these varieties are closer to Calabi-Yau then to $\mathbb{P}^n$, thus their rigidity is easier to prove in higher dimension.

**Theorem 2.10** (Inversion of adjunction, [11, Theorem 17.7])**.** *Let $X$ be a normal variety and let $S \subset X$ be an irreducible divisor with singularities of codimension $\geqslant 3$. Let $B$ be an effective $\mathbb{Q}$-Cartier divisor. Assume that $K_X + S$ is $\mathbb{Q}$-Cartier and is terminal. Then $K_X + S + B$ is log canonical in a neighborhood of $S$ if and only if $K_S + B$ is log canonical.*

We use the inversion of adjunction to study singularities of linear systems as follows. Let $F$ be a prime divisor on a variety $X$ and suppose there is a divisor $D$ such that $K_X + D$ is not canonical along $B \subset F$. Suppose that $B$ is an ordinary double point of $F$ and $\dim X \geqslant 4$ or that $F$ is smooth along $B$. Then $K_X + D + F$ is not log canonical along $B$ and by adjunction $K_F + D\big|_F$ is not log canonical.

**Theorem 2.11** ([3, Theorem 1.5])**.** *Let $Z$ be a smooth curve. Suppose that there is a commutative diagram*

$$\begin{array}{ccc} V & \overset{\rho}{\dashrightarrow} & V' \\ \downarrow{\scriptstyle \pi} & & \downarrow{\scriptstyle \bar{\pi}} \\ Z & = & Z \end{array}$$

*such that $\pi$ and $\bar{\pi}$ are flat morphisms, and $\rho$ is a birational map that induces an isomorphism*

$$\rho\big|_{V\setminus X} : V \setminus X \to \bar{V} \setminus \bar{X}$$

*where $X$ and $\bar{X}$ are scheme fibers of $\pi$ and $\bar{\pi}$ over a point $O \in Z$, respectively. Suppose that the varieties $V$ and $\bar{V}$ have terminal $\mathbb{Q}$-factorial singularities, the divisors $-K_V$ and $-K_{\bar{V}}$ are $\pi$-ample and $\bar{\pi}$-ample respectively, the fibers $X$ and $\bar{X}$ are irreducible, the variety $X$ has log terminal singularities, and $\alpha(X) \geqslant 1$. Then $\rho$ is an isomorphism.*

*Proof taken from [3].* Suppose that the variety $X$ has log terminal singularities, the inequality $\alpha(X) \geqslant 1$ holds, and $\rho$ is not an isomorphism. Let $D$ be a generic very ample divisor on $Z$. Put

$$\Lambda = \big|-nK_V + \pi^*(nD)\big|,\ \Gamma = \big|-nK_{\bar{V}} + \bar{\pi}^*(nD)\big|,\ \bar{\Lambda} = \rho(\Lambda),\ \bar{\Gamma} = \rho^{-1}(\Gamma),$$

where $n$ is a natural number such that $\Lambda$ and $\Gamma$ have no base points. Put

$$M_V = \frac{2\varepsilon}{n}\,\Lambda + \frac{1-\varepsilon}{n}\,\overline{\Gamma},\ M_{\bar{V}} = \frac{2\varepsilon}{n}\,\bar{\Lambda} + \frac{1-\varepsilon}{n}\,\Gamma,$$

where $\varepsilon$ is a positive rational number.

The log pairs $(V, M_V)$ and $(\bar{V}, M_{\bar{V}})$ are birationally equivalent, and for $\varepsilon$ small enough $K_V + M_V$ and $K_{\bar{V}} + M_{\bar{V}}$ are ample. We claim that the map $\rho$ is biregular if the singularities of both log pairs $(V, M_V)$ and $(V, M_{\bar{V}})$ are canonical. Consider the resolution $(V', M')$ of the map $\rho$, we can write

$$K_{V'} + M' \sim_{\mathbb{Q}} K_V + M_V + \sum a_i E_i \sim_{\mathbb{Q}} K_{\bar{V}} + M_{\bar{V}} + \sum b_i E_i.$$

Since the pairs are canonical the coefficients $a_i$ and $b_i$ are non-negative for all $i$. Thus, applying the negativity lemma we see that the maps from $V'$ to $V$ and $\bar{V}$ have the same exceptional divisors. Hence $\rho$ is an isomorphism in codimension 1 and by Proposition 2.6 it is an isomorphism.

The linear system $\Gamma$ does not have base points. Thus, there is a rational number $\varepsilon$ such that the log pair $(\bar{V}, M_{\bar{V}})$ is canonical. So the log pair $(V, M_V)$ is not canonical. The log pair

$$\left(V,\ X + \frac{1-\varepsilon}{n}\,\bar{\Gamma}\right)$$

is not log canonical, because $\Lambda$ does not have not base points, and $\bar{\Gamma}$ does not have base points outside of the fiber $X$, which is a Cartier divisor on the variety $V$. The log pair

$$\left(X,\ \frac{1-\varepsilon}{n}\,\bar{\Gamma}\big|_X\right)$$

is not log canonical by inversion of adjunction, which is impossible because $\alpha(X) \geqslant 1$. □

Let us formulate the converse to the Theorem 2.9: if there exists fiber $F$ with $\alpha(F) < 1$, then there exists non-trivial fiberwise map $V \dashrightarrow V'$. This statement does not hold: variety $X$ from the Theorem 1.1 is an counterexample. It is interesting to know the answer to the following question.

**Question 2.** *What conditions on a fibration and a fiber imply the existence of a non-trivial fiberwise map?*

# 3 Example in dimension $\geqslant$4

In this section we construct examples in dimension $\geqslant 4$ and prove part (i) of Theorem 1.1.

Let $W$ be a generic divisor of degree $(2M, 2l)$ on $\mathbb{P}^M \times \mathbb{P}^1$, $M \geqslant 3$, $l \geqslant 3$. Let $g : V \to \mathbb{P}^M \times \mathbb{P}^1$ be a double cover ramified over $W$. Because $W$ is generic (in particular smooth), variety $V$ is smooth. Let $\pi : V \to \mathbb{P}^1$ be a composition of $g$ and projection onto $\mathbb{P}^1$. The fiber of $\pi$ is a double cover of $\mathbb{P}^M$ ramified over a hypersurface of degree $2M$. Since $W$ is generic we may also assume that every fiber is smooth or has one ordinary double point and $\mathrm{Pic}(F) = K_F \cdot \mathbb{Z}$. Hence a generic fiber $F$ is a smooth Fano variety. Therefore $\mathrm{Pic}(V) = \mathbb{Z} \oplus \mathbb{Z}$ and $V$ is a Fano fibration over projective line.

We provide the specific case of the following theorem for the sake of simplicity.

**Theorem 3.1** ([19, Theorem 2.1])**.** *Let $\pi : V \to \mathbb{P}^1$ be a Fano fibration, suppose $\dim V \geqslant 4$ and $\mathrm{Pic}(V) = K_V \cdot \mathbb{Z} \oplus F \cdot \mathbb{Z}$, where $F$ is a fiber of $\pi$. Suppose $V$ satisifies the $K$-condition, that is, $\left| -nK_V \right|$ is not movable for*

*any $n$. Suppose the generic fiber of $\pi$ is birationally superrigid. Assume that $\alpha(F) \geqslant 1$ and $\mathrm{Sing}(F)$ consists only of ordinary double points for every fiber $F$ of $\pi$. Then $V$ is birationally superrigid.*

The nature of the $K$-condition is the following. For superrigidity is easier to prove for Fano varieties which are close to Calabi-Yau varieties. In case of fibrations over projective line we do want for anticanonical class to be small on fibers, that is that fibers are similar enough to Calabi-Yau, but there is additional requirement. Suppose $\operatorname{Pic} X = \mathbb{Z} \cdot D \oplus \mathbb{Z} \cdot F$, where $F$ is a fiber, and define $a$ and $b$ by the $K_X = aD + bF$. Condition of being small on the fibers is $a = -1$. If $b < 0$ or is just small enough, then $X$ is a Fano variety and thus has more than one Mori fiber space structure. The $K$-condition is that $b$ is big enough. Thus not only we want $-K_X$ to be small enough, we want $K_X$ to be positive in some sense.

*Proof.* Let $f$ be a birational map from $V$ to a Mori fiber space $X$ and suppose $f$ is not fiberwise. Consider a linear system $\mathcal{M}_X$ which defines the Fano fibration $g : X \to S$. Let $\mathcal{M} = f^{-1}\mathcal{M}_X$ be the proper transform of $\mathcal{M}_X$, then

$$\mathcal{M} \subset \big| -nK_V + sF \big|. \tag{2}$$

If $n = 0$ then $f$ is clearly fiberwise, hence $n > 0$. Since $\mathcal{M}$ is a movable system it follows from the $K$-condition that $s > 0$. We can rewrite (2) as

$$K_V + \lambda\mathcal{M} \sim_{\mathbb{Q}} lF, \tag{3}$$

where $l > 0$ because $s > 0$. Consider the resolution $W$ of the birational map $f$. Let $\psi : W \to V$ and $\varphi : W \to X$ be the birational morphisms corresponding to the resolution. Denote $\mathcal{M}_W = \varphi^{-1}\mathcal{M}$, then we can write

$$\psi^*(K_V + \lambda\mathcal{M}) + \sum a_i E_i \sim_{\mathbb{Q}} K_W + \mathcal{M}_W \sim_{\mathbb{Q}} \varphi^*(K_X + \lambda\mathcal{M}_X) + \sum b_j E_j,$$

where $E_i$ and $E_j$ are exceptional divisors of $\psi$ and $\varphi$ respectively. Intersecting this equality with a generic divisor $D \in \varphi^*\mathcal{M}$ and applying (2) we get

$$D \cdot \psi^*(lF) + \sum a_i D \cdot E_i \sim_{\mathbb{Q}} D \cdot \varphi^*(K_X).$$

Since $K_X|_D$ is anti-ample there is a coefficient $a_i < 0$ for some $i$, therefore $(V, \lambda D)$ is not a canonical pair. Since this is true for a generic divisor $D$, the pair $(V, \lambda\mathcal{M})$ is not canonical. Denote the centre of this non-canonical singularity by $Z = \psi(E_i)$. Suppose that $Z$ does not lie on any fiber of $\pi$ and consider a generic fiber $F$. The pair $(F, \lambda\mathcal{M}|_F)$ is not canonical at $Z|_F$ and

$$\lambda\mathcal{M} \sim_{\mathbb{Q}} (-K_V + lF)|_F = -K_F,$$

which contradicts the superrigidity of a generic fiber $F$ by Proposition 2.7. Thus $Z$ lies on some fiber $F$. The pair $(V, F)$ is terminal since $\mathrm{Sing}(F)$ consists of ordinary double points or is empty, thus we can apply Inversion of adjunction. Hence the pair $(F, \mathcal{M}|_F)$ is not log canonical at $Z$ which contradicts the bound on the $\alpha$-invariant. □

**Corollary 3.2.** *Let $\pi : V \to \mathbb{P}^1$ be a Fano fibration, suppose $\dim V \geqslant 4$ and $\mathrm{Pic}(V) = K_V \cdot \mathbb{Z} \oplus F \cdot \mathbb{Z}$, where $F$ is a fiber of $\pi$. Suppose $V$ satisfies $K$-condition. Suppose the generic fiber of $\pi$ is birationally superrigid and assume that $\alpha(F) \geqslant 1$ for every fiber $F$ of $\pi$. Then all birational maps from $V$ to any Mori fiber space are automorphisms $V \to V$.*

*Proof.* By Theorem 3.1 the variety $V$ is birationally superrigid, therefore every map from $V$ to a Mori fiber space is a fiberwise map. By Theorem 2.9 every fiberwise map is an isomorphism. □

Here we describe the sufficient conditions (slight modification of those in [17]) for the fiber to be superrigid and to have $\alpha(F) > 1$. Then we show that most double covers of $\mathbb{P}^M$ ramified over hypersurface of degree $2M$ satisfy these conditions and therefore every fiber of a generic $V$ satisfies these conditions as well.

Let $W \subset \mathbb{P}^M$ be a hypersurface of degree $2M$, $M \geqslant 3$. For a nonsingular point $x \in W$ fix a system of affine coordinates $z_1, \ldots, z_M$ on $\mathbb{P}^M$ with origin at $x$ and set

$$w = q_1 + q_2 + \cdots + q_{2M}$$

to be the equation of the hypersurface $W$, where $q_i = q_i(z)$ are homogeneous polynomials of degree $\deg q_i = i$. We can assume that $q_1 = z_1$. Denote

$$\bar{q}_i = \bar{q}_i(z_2, \ldots, z_M) = q_i\big|_{\{z_1=0\}} = q_i(0, z_2, \ldots, z_M).$$

Suppose $M \geqslant 5$. We say that $W$ satisfies the regularity conditions at a smooth point $x$ if the rank of the quadratic form $\bar{q}_2$ is at least 2.

For $M = 4$ we need either

(i) The rank of the quadratic form form $\bar{q}_2$ is at least 2,

(ii) or rank of $\bar{q}_2$ is 1 and the following additional condition is satisfied. Without loss of generality we assume in this case that $\bar{q}_2 = z_2^2$. Now we require that the the polynomials $\bar{q}_3(0, z_3, 0) = q_3(0, 0, z_3, 0)$ and $\bar{q}_4(0, z_3, 0)$ are not zero.

Suppose $M = 3$. Then we require either

(i) The rank of the quadratic form form $\bar{q}_2$ is at least 2,

(ii) rank of $\bar{q}_2$ is 1 and at least one of the following polynomials in the variable $z_3$,

$$\bar{q}_3(0, z_3), \quad \bar{q}_4(0, z_3), \quad \textit{or} \quad \bar{q}_5(0, z_3)$$

is not zero,

(iii) or $\bar{q}_2 = 0$ and polynomial $\bar{q}_3(1, t)$ has 3 distinct roots.

Let $W \subset \mathbb{P}^M$ be a hypersurface of degree $2M$ satisfying the regularity conditions at every nonsingular point. Suppose $\mathrm{Sing}(W)$ is empty or consists of only ordinary double points, then we call $W$ a regular hypersurface. Let $\sigma : F \to \mathbb{P}^M$, $M \geqslant 3$, be the double cover ramified over a regular hypersurface $W \subset \mathbb{P}^M$ then we say $F$ is a regular double space.

**Proposition 3.3** ([18, Subsection 1.3])**.** *Let $X$ be a regular double space. Suppose $D \sim -nK_X$ is a Cartier divisor on $X$. Let $o$ be an ordinary double point on $X$. Then $(X, \frac{1}{n}D)$ is canonical at $o$.*

A regular double space $X$ of dimension $M$ can be embedded into $\mathbb{P}(1, \ldots, 1, M)$ as a hypersurface of degree $2M$ which does not pass through singular point, therefore there is a natural definition of multiplicity of subvarieties of $X$ at a point.

The following theorem is a stronger version of [18, Theorem 4].

**Theorem 3.4.** *Let $X$ be a regular double space. Then for every effective $\mathbb{Q}$-divisor $D$ such that $D + K_X \sim_{\mathbb{Q}} 0$ the pair $(X, D)$ is canonical.*

*Proof.* Note that if the pairs $(X, D)$ and $(X, D')$ are canonical, then the pair $(X, \alpha D + (1 - \alpha)D')$ is also canonical for $0 \leqslant \alpha \leqslant 1$. Thus without the loss of generality we may assume that $D$ is irreducible. Suppose $D \sim -nK_X$ and that the pair $(X, \frac{1}{n}D)$ is not canonical.

It follows from the Proposition 3.3 that singular points cannot be non-canonical centers of $(X, \frac{1}{n}D)$. It was proven in [17, Lemma 5] that the pair $(X, \frac{1}{n}D)$ is canonical at curves and at nonsingular points do not lie on the ramification divisor. Thus $(X, \frac{1}{n}D)$ is not canonical at a nonsingular point $x \in X$ on the ramification divisor and $\mathrm{mult}_x D > 1$. Let $A$ be a hyperplane in $\mathbb{P}^M$ tangent to $W$ at the point $z = \sigma(x)$. Denote $\sigma^{-1}(A)$ as $H$. Note that $H$ is irreducible, $\mathrm{mult}_x H \geqslant 2$, and $H \sim -K_X$ is a generator of $\mathrm{Pic}(X)$. If $D \neq H$, then for the cycle $Z = H \cdot D$ we get

$$\deg Z = 2n, \quad \mathrm{mult}_x Z \geqslant 2\,\mathrm{mult}_x D > 2n.$$

As there is no such cycle $Z$ we conclude that $D = H$. Hypersurface $H$ is a double cover of $A$ ramified over $W_A = W \cap A$. Since $W$ is non-singular the point $x$ is an isolated singularity of $W_A$. Thus $H$ has isolated singularity at $x$ and local equation of $H$ is

$$y^2 = \bar{q}_2(z_2, \dots, z_M) + \bar{q}_3(z_2, \dots, z_M) + \dots + \bar{q}_{2M}(z_2, \dots, z_M).$$

Claim: if $\operatorname{rank} \bar{q}_2 > 1$ then the pair $(X, H)$ is canonical. Let $f : X^+ \to X$ be the blow up of $x$, $E$ be an exceptional divisor and $H^+$ be a proper transform of $H$. Then

$$K_{X^+} + H^+(3 - M)E = f^*(K_X + H) = 0,$$

also recall that $M \geqslant 3$. The pair $(X^+, H^+(3 - M)E)$ is non-canonical, hence so is $(X^+, H^+)$. Denote $H^+|_E$ as $Q$. By adjunction the pair $(E, Q)$ is not log canonical. Hypersurface $Q \in P^{M-1} = E$ is the quadric given by $y^2 = \bar{q}_2(z_2, \dots, z_M)$. Note that $Q$ is singular, otherwise the pair $(E, Q)$ is canonical. Since the rank of $\bar{q}_2$ is more than 1 the a singular locus $S$ of $Q$ is of codimension 3 in $E$, in particular $Q$ is irreducible. For a blow up $g : E^+ \to E$ of $S$ we can write

$$K_{E^+} + Q^+(3 - \operatorname{codim}_E S)F = g^*(K_E + Q),$$

where $F$ is an exceptional divisor. As $Q^+$ is smooth the pair $(E^+, Q^+)$ is log canonical. Since coefficient at $R$ is non-positive, the pair $(E, Q)$ is log canonical, hence $(X, H)$ is canonical.

In particular this implies that if $M = 5$ then $(X, H)$ is canonical.

Suppose $M = 4$ and $\operatorname{rank} \bar{q}_2 = 1$, then $H$ has equation

$$y^2 = z_2^2 + z_2(\dots) + c_1 z_3^3 + c_2 z_3^4 + z_4(\dots) + \dots$$

As $c_1$ or $c_2$ is not zero, $H$ has singularity at worst of the type $cA_3$ at $x$. Thus $H$ is terminal and $(X, H)$ is canonical.

Now suppose $M = 3$ and $\bar{q}_2 = z_2^2$. It follows from the regularity conditions that equation of $H$ is $y^2 = z_2^2 + z_3^3$, $y^2 = z_2^2 + z_3^4$, $y^2 = z_2^2 + z_3^5$ or $y^2 = z_2(z_2^2 + z_3^2)$ thus $x$ is a Du Val singularity, $H$ is canonical, and therefore the pair $(X, H)$ is canonical. □

By combining Proposition 2.7 and this theorem we obtain the following statement.

**Corollary 3.5.** *Let $F$ be a regular double space of dimension $\geqslant 3$. Then $F$ is superrigid and in particular $\alpha(F) \geqslant 1$.*

It is proven in [17, Proposition 5] that a generic hypersurface of degree $2M$ is regular. Thus a generic fiber of a generic fibration onto regular double spaces is regular. We want every fiber to be regular.

**Proposition 3.6.** *Let $\mathcal{W}$ be the space of all hypersurfaces of degree $2M$ in $\mathbb{P}^M$. Denote the space of all regular hypersurfaces by $\mathcal{W}_{reg} \subset \mathcal{W}$. If $M \geqslant 3$ then the codimension of $\mathcal{W} \backslash \mathcal{W}_{reg}$ is 2 or more.*

*Proof.* First note, that codimension of hypersurfaces with $\geqslant 2$ ordinary double point or with worse singularities is 2. Thus it is enough to prove that the space of hypersurfaces which do not satisfy the regularity conditions at nonsingular points has codimension $\geqslant 2$. Consider the variety $V$ and the incidence hypersurface $I$ in it defined by

$$V = \mathbb{P}^M \times H^0(\mathbb{P}^M, \mathcal{O}(2M)),$$
$$I = \{(x, F) \in V \mid F(x) = 0\}.$$

Let $p$ and $q$ be natural projections $p : I \to \mathbb{P}^M$ and $q : I \to H^0(\mathbb{P}^M, \mathcal{O}(2M))$. Let $Y$ be the subset of "bad" pairs

$$Y = \{(x, F) \in I \mid \text{F is not regular at } x \text{ if } x \text{ is smooth,} \\ \text{or } x \text{ is worse than an ordinary double point}\}.$$

To prove the proposition it is enough to show that $\operatorname{codim} q(Y) \geqslant 2$. To show this it is sufficient to prove that $\operatorname{codim}_{p^{-1}(x)} p^{-1}(x) \cap Y \geqslant M + 1$.

Consider the equation of $W$ in affine coordinates in the neighborhood of a point $x$

$$q_1 + q_2 + \cdots + q_{2M} = 0$$

If $q_1 = 0$ then $W$ is singular at $x$. If $q_2$ is non-degenerate then $x$ is an ordinary double point point of $W$. Thus, the locus of hypersurfaces of degree $2M$ for which $x$ is not an ordinary double point is of codimension $M + 1$ in $p^{-1}(x)$.

If $q_1 \neq 0$, we may assume that $q_1 = z_1$ and take $\bar{q_2} = q_2(0, z_2, \ldots, z_M)$. The set of quadratic forms of rank $\leqslant 1$ in the variables $z_2, \ldots, z_M$ is of codimension

$$c(M) = \frac{(M-1)(M-2)}{2}.$$

When $M \geqslant 5$ we have $c(M) \geqslant M + 1$. Suppose $M = 4$. As conditions $\bar{q_3}(0, z_3, 0) = 0$ and $\bar{q_4}(0, z_3, 0) = 0$ add 2 to the codimension, the codimension is $c(4) + 2 = 5$. If $M = 3$ than we need add 4 to the codimension as $c(3) = 1$.

Similarly to the previous case varieties with $\operatorname{rank}(\bar{q}_2) = 2$ satisfying one of the conditions: $\bar{q}_3(0, z_3) = 0$, $\bar{q}_4(0, z_3) = 0$ and $\bar{q}_5(0, z_3) = 0$, are of codimension 4. The set of varieties with $\bar{q}_2 = 0$ is of codimension 3 and condition: $\bar{q}_3(1, t) = 0$ does not have 3 distinct roots, adds 1 to the codimension. □

**Lemma 3.7.** *Suppose $V$ is a variety from Theorem 1.1 (i). Then the anti-canonical class of the variety $V$ is not big and $V$ satisfies $K$-condition.*

*Proof.* It is easy to compute that

$$-K_V = g^*(L),$$

where $L$ is a divisor of bidegree $(1, 2-l)$, $l \geqslant 3$. Thus the $K$-condition holds and $-K_V$ is not big. □

*Proof of Theorem 1.1 (i).* Let $g : V \to \mathbb{P}^1$ be the map giving the structure of Fano fibration on $V$. We can view $g$ as a map from $\mathbb{P}^1$ to the space of hypersurfaces of degree $2M$ in $\mathbb{P}^M$. By Proposition 3.7, the codimension of the set of non regular hypersurfaces is 2 and therefore a generic rational curve of degree $2l$ does not intersect this set. Thus for a generic divisor $W$ every fiber of $g$ is regular. The variety $V$ satisfies $K$-condition by Lemma 3.8, hence Theorem 3.5 implies that a generic fiber is birationally rigid and $\alpha(F) \geqslant 1$ for every fiber $F$.

Hence, $V$ satisfies conditions of the Corollary 3.2 and we conclude that any birational map from $V$ to a Mori fiber space is an isomorphism. □

# 4 Example in dimension 3

In this section we construct an example in dimension 3 and prove part $(ii)$ of Theorem 1.1.

Let $Y = (\mathbb{C}^6 \setminus Z\big(\langle u, v\rangle \cap \langle x, y, z, w\rangle\big)/(\mathbb{C}^*)^2$, where $(\mathbb{C}^*)^2$-action is given by a matrix

$$\begin{pmatrix} u & v & x & y & z & w \\ 0 & 0 & 1 & 1 & 2 & 3 \\ 1 & 1 & -3 & -3 & 0 & 0 \end{pmatrix}$$

and let

$$Q = w^2 + z^3 + (u^{12} + v^{12})M_4(x, y)z + R_{18}(u, v)x^2y^2(x - y)^2,$$

where $M_4 \in \mathbb{C}[x, y]$, $\deg M_4 = 4$ and $R_{18} \in \mathbb{C}[u, v]$, $\deg R_{18} = 18$. Let $\mathcal{L}$ be the set of hypersurfaces given by $Q = 0$. Clearly $\mathcal{L}$ is a linear system of

divisors. Let $X$ be a generic element of $\mathcal{L}$. The variety $Y$ is a projective simplicial toric variety with the Cox ring $\mathbb{C}[u, v, x, y, z, w]$. This ring has 2 gradings given by the matrix above, for example $X$ is a hypersurface is of degree $(6, 0)$.

A fiber of a map $\pi : Y \to \mathbb{P}^1$ defined by

$$(u : v : x : y : z : w) \mapsto (u : v)$$

restricted to $X \in \mathcal{L}$ is del Pezzo surface of degree 1. Indeed, for a fixed $(u : v)$ the fiber is a hypersurface of degree 6 in $\mathbb{P}(1, 1, 2, 3)$ given by $w^2 = z^3 + z^2P_2(x, y) + zP_4(x, y) + P_6(x, y)$, where $P_i$ is a homogeneous polynomial of degree $i$. Thus each fiber is a double cover of a quadratic cone $\mathbb{P}(1, 1, 2)$ in $\mathbb{P}^3$ ramified over a cubical section $z^3 + z^2P_2(x, y) + zP_4(x, y) + P_6(x, y) = 0$.

Denote $Z = \{u, v, x, y, z, w\}$. Denote the torus-invariant divisor given by $s = 0$, $s \in Z$ as $D_s$, and note that $D_u \sim D_v \sim F$, where $F$ is a fiber of $\pi$. It is easy to see that $\mathrm{Cl}(Y) = \mathbb{Z} \cdot D_y \oplus \mathbb{Z} \cdot F$.

**Lemma 4.1.** *Let $F$ be a fiber of $\pi : Y \to \mathbb{P}^1$ and suppose $X \in \mathcal{L}$. Then*

*(i) $X$ is a Cartier divisor*

*(ii) $\big|X\big| = \big|6D_y + 18F\big|$ is base point free,*

*(iii) $\big|X + F\big|$ is ample,*

*(iv) $\big|X\big|$ is big.*

*Proof.* The divisor $X$ is Cartier because it does not pass through a singular locus of $Y$, that is curves $x = y = z = 0$ and $x = y = w = 0$ because coefficients at monomials $w^2$ and $z^3$ are never 0.

The equation of $D \in \big|X\big|$ may contain monomials: $w^2$, $z^3$, $x^6u^{18}$, $x^6v^{18}$, $y^6u^{18}$ and $y^6v^{18}$ which are not equal zero at the same time at any point on $Y$, thus $(ii)$ holds.

We claim that $X+F$ is ample by the Nakai-Moishezon criterion. Indeed, the system $\big|X+F\big|$ is obviously base point free, therefore it is enough to check intersections with curves. Suppose $C$ is a curve in a fiber then

$$C \cdot (X + F) = C \cdot X > 0.$$

If $C$ is not in a fiber then by $(ii)$ $C \cdot (X + F) \geqslant C \cdot F > 0$.

Clearly $2X \sim X + F + (6D_y + 17F)$, hence $(iv)$ follows from $(iii)$. □

**Lemma 4.2.** *Let $X \in \mathcal{L}$ be a generic member of the family. Then the following assertions hold:*

(i) *There are* 108 *cuspidal curves of anticanonical degree* 1 *in fibers of* $X$, 72 *of them are given by equations* $R_{18} = M_4 = 0$. *Other* 36 *curves are given by* $u^{12} + v^{12} = 0$ *and* $x = 0$, $y = 0$ *or* $x = y$.

(ii) *Let* $F$ *be a fiber and let* $C \in F$ *be one of the* 72 *curves. Let* $P$ *be the cusp of* $C$, *then* $P$ *is an ordinary double point of* $F$.

(iii) *Let* $C$ *be one of* 36 *curves and let* $P$ *be the cusp of* $C$. *Then* $P$ *is an ordinary double point of* $X$.

(iv) *The variety* $X$ *is smooth outside of* 36 *ordinary double points described in* (*iii*).

*Proof.* The coefficient at $z^2$ is zero for every fiber, therefore the equations

$$R_{18}x^2y^2(x-y)^2 = 0$$
$$(u^{12} + v^{12})M_4(x, y) = 0$$

give us all cuspidal curves of degree 1. As $R_{18}$ and $M_4$ are generic we must have $R_{18} = M_4 = 0$ or $u^{12} + v^{12} = x^2y^2(x-y)^2 = 0$, moreover, $M_4$ and $R_{18}$ do not have multiple roots. Thus $(i)$ holds. Note that the cusps of these curves are at $w = z = 0$.

The local equation of $X$ at the cusp of one of the 72 curves is

$$w^2 + z^3 + zs + t = 0.$$

Clearly the fiber $t = 0$ has an ordinary double point.

If $C$ is one of the 36 curves, then the local equation is

$$w^2 + z^3 + zt + s^2 = 0,$$

thus $X$ has an ordinary double point.

Note that varieties from $\mathcal{L}$ do not pass through singular locus of $Y$. Obviously $w = 0$ at any singular point. Let $P \in Y$ be a point such that $xy(x-y) \neq 0$. There is a linear function $t(u, v)$ such that $t \neq 0$ at $P$. If $P$ does not lie on $X_R$ corresponding to $R = t^{18}$, then by Bertini's theorem [13, Theorem 4.1] generic $X$ is smooth at $P$. If $P$ lies on $X_R$ then we can replace $t^{18}$ with $ct^{18}$ for some $c \in \mathbb{C}$ to get $X_R$ which does not go through $P$. Therefore $x = 0$, $y = 0$ or $x = y$.

Suppose $zQ_{12} \neq 0$ at $P$. Then repeating the argument for some linear function $s(x, y)$ and $P = s^4$ we get that generic $X$ is smooth at $P$ by Bertini's theorem. Thus $z = 0$ or $Q_{12} = 0$. Notice that $Q_{12} = 0$ if and only if $z = 0$, because $M_4$ does not have common roots with $xy(x-y)$ as it is generic and $\frac{\partial Q_P}{\partial z} = 3z^2 + Q_{12}M_4 = 0$.

Thus every singular point must satisfy $w = z = Q_{12} = xy(x-y) = 0$. There are 36 points satisfying these equations and they are described in (*iii*). □

Now we will prove that $X$ is a Mori fiber space. We already know that it is terminal, thus we need to show that it is $\mathbb{Q}$-factorial and that relative Picard rank $\rho(X/\mathbb{P}^1) = 1$. We intend to use the following theorem.

**Theorem 4.3** ([20, Theorem 4.1])**.** *Let $Y$ be a complete simplicial toric fourfold and let $X \subset Y$ be a hypersurface such that $\mathrm{Sing}(Y) \cap X = \emptyset$ and $\mathrm{Sing}(X)$ consists only of ordinary double points. Let $\widetilde{X}$ be a resolution of $X$ obtained by blow ups of all the ordinary double points. Let $\mu_X$ be the number of singular points on $X$ and let $\mathcal{I}_{\mathrm{Sing}(X)}$ be the sheaf of ideals on $Y$ of singular points of $X$. Assume that the following conditions are satisfied:*

*(i)* $H^i(Y, \mathcal{O}_Y(-2X)) = 0$ *for* $i = 1, 2, 3$*;*

*(ii)* $H^i(Y, \mathcal{O}_Y(-X)) = 0$ *for* $i = 1, 2, 3$*;*

*(iii)* $H^i(Y, \bar{\Omega}^1_Y \otimes \mathcal{O}_Y(-X)) = 0$ *for* $i = 1, 2, 3$*;*

*(iv)* $\delta_X = H^0(Y, \mathcal{O}(K_Y + 2X) \otimes \mathcal{I}_{\mathrm{Sing}(X)}) - (H^0(Y, \mathcal{O}(K_Y + 2X)) - \mu_Y) = 0$

*Then* $h^{1,1}(\widetilde{X}) = h^1(\bar{\Omega}^1_Y) + \mu_X$.

The number $\delta_Y$ is the difference between expected and actual dimension of the space and is always non-negative. It is positive when the singular points are not in a generic position. It is also known as a defect [20].

**Lemma 4.4.** *For a generic $X \in \mathcal{L}$ the defect $\delta_X = 0$.*

*Proof.* Let $P_i$, $i = 1 \dots 36$ be the ordinary double points of $X$. It is enough show that there are divisors $H_i \in \left|2X + K_Y\right| = \left|5D_x + 19F\right|$, $i = 1 \dots 36$ such that $H_i$ passes through $P_j$ for any $j \neq i$ and does not pass through $P_i$. Let $F_k$, $k = 1 \dots 12$ be the fibers containing double points of $X$. Let $D_{x-y}$ be the divisor defined by $y = x$. Taking 11 fibers $F_k$ and 2 out of $D_x$, $D_y$, $D_{x-y}$ we get a divisor $H'_i$ linearly equivalent to $2D_x + 11F$ passing through any 35 singular points out of 36. The divisor $A$ given by the equation $(x - 2y)^3 u^8$ does not pass through any singular points of $X$, therefore $H_i = H'_i + A$. □

**Proposition 4.5.** *Suppose $X \in \mathcal{L}$ is generic. Then $X$ is $\mathbb{Q}$-factorial and $\mathrm{Pic}\, X \otimes \mathbb{Q} = \mathbb{Q} \cdot D_y|_X \oplus \mathbb{Q} \cdot F$, where $F$ is a fiber of $\pi|_X$.*

*Proof.* Let us show that $X$ satisfies the conditions of Theorem 4.3. By Lemma 4.1 the linear system $\big|X\big|$ is big and nef, therefore it follows from the toric Kawamata-Viehweg vanishing [4, Theorem 9.3.10] that $H^i(Y, \mathcal{O}_Y(K_Y + 2X)) = 0$. Thus by the toric Serre Duality [4, Theorem 9.2.10] we get $(i)$. By the same argument we get $(ii)$.

As $\mathrm{Cl}(Y) = \mathbb{Z} \oplus \mathbb{Z}$ the following sequence is exact by [4, Theorem 8.1.6]

$$0 \to \bar{\Omega}^1_Y(-X) \to \bigoplus_{s \in Z} \mathcal{O}_Y(-D_s - X) \to \mathcal{O}_Y(-X) \oplus \mathcal{O}_Y(-X) \to 0.$$

By taking associated long exact sequence of cohomologies and by using $(ii)$ we get

$$0 \to H^i(Y, \bar{\Omega}^1_Y(-X)) \to \bigoplus_{s \in Z} H^i(Y, \mathcal{O}_Y(-D_s - X)), \quad i = 1, 2, 3.$$

It is enough to prove that $H^i(Y, \mathcal{O}_Y(-D_s - X)) = 0$ for all $s \in Z$, $i = 1, 2, 3$. If $s \in u, v, z, w$ then $D_s + X$ is big and nef, therefore by the Serre duality and the Kawamata-Viehweg vanishing $H^i(Y, \mathcal{O}_Y(-D_s - X)) = 0$ holds. Let $F$ be a fiber of $\pi$. Then $\mathcal{O}_F(F) = \mathcal{O}_F$ and the sequence

$$0 \to \mathcal{O}_Y \to \mathcal{O}_Y(F) \to \mathcal{O}_F \to 0$$

is exact. Hence the sequences

$$H^i(Y, \mathcal{O}_Y) \to H^i(Y, \mathcal{O}_Y(F)) \to H^i(F, \mathcal{O}_F), \quad i = 1, 2, 3$$

are exact. Applying [4, Theorem 9.3.2] again we get $H^i(Y, \mathcal{O}_Y(F)) = 0$ from these sequences. As $-X - D_y \sim -X - D_y \sim K_Y - F$ by the Serre duality

$$H^i(Y, \mathcal{O}_Y(-D_x - X)) = H^{4-i}(Y, \mathcal{O}_Y(F)) = 0.$$

Thus by Theorem 4.3 $\operatorname{rk} \mathrm{Pic}(\widetilde{X}) = 38$. On the other hand $\operatorname{rk} \mathrm{Pic}(\widetilde{X}) = \operatorname{rk} \mathrm{Cl}(X) + 36$, thus $\operatorname{rk} \mathrm{Pic}(X) = \operatorname{rk} \mathrm{Cl}(X) = 2$ and $X$ is $\mathbb{Q}$-factorial. Clearly, divisors $D_y|_X$ and $F|_X$ are the generators. □

Suppose we have a non-regular birational map $f$ from $X$ to a Mori fiber space and suppose it is given by a linear system $\mathcal{M}$. Suppose that $f$ is not regular at $Y \subset F \subset X$, where $F$ is a fiber of $\pi : X \to \mathbb{P}^1$, that is $Y \subset$ is in base locus of $(\mathcal{M})$. Because $f$ is a map to a Mori fiber space it can be shown, that $\mathcal{M}$ is very singular in some subset of $Y$. Corollary 4.8 implies that this subset must be a cusp of a degree 1 curve in $F$. On the other hand we show that $\mathcal{M}$ cannot be very singular at ordinary double points of $X$ or of fibers of $\pi$. By Lemma 4.2 all cusps of degree 1 curves in fibers are also ordinary double points of $X$ or fibers of $\pi$.

**Lemma 4.6.** *The variety $X$ satisfies $K$-condition and $-K_X$ is not big.*

*Proof.* It is easy to see that $K_Y = -7D_y - 17F$, thus by the adjunction the anticanonical class of $X$ is $-K_X = D_y|_X - F|_X$. Since $\left| -nK_X \right| = \emptyset$ the system is not movable for any $n$ and $-K_X$ is not big. □

**Lemma 4.7** ([10, Proposition 3.2])**.** *Let $F$ be a del Pezzo surface of degree 1 and let $C \in \left| -K_F \right|$ be an irreducible curve. Suppose $(F, C)$ is not log canonical at $P$ and $F$ is smooth at $P$. Then $C$ is a cuspidal rational curve.*

**Corollary 4.8.** *Let $F$ be a del Pezzo surface of degree 1 with at worst ordinary double points and let $D \sim_{\mathbb{Q}} -K_F$ be a $\mathbb{Q}$-Cartier divisor. If $(F, D)$ is not log canonical at $P$, then there is a rational curve $C \in \left| -K_F \right|$ such that $P \in C$ is a cusp.*

**Lemma 4.9** ([6, Theorem 3.10])**.** *Let $X$ be a 3-dimensional variety and let $D$ be an effective $\mathbb{Q}$-Cartier divisor on $X$ such that the pair $(X, D)$ is not canonical and a point $P \in X$ is its center. Suppose $P$ is an ordinary double point on $X$ and let $f : X^+ \to X$ be a blow up of $P$ with the exceptional divisor $E$. Then $a(E, X, D) < 0$.*

*Proof of Theorem 1.1 (ii).* Suppose there is such map. Then, similarly to the proofs of Proposition 2.7 and Theorem 3.1, there is a mobile linear system $\mathcal{M} \in \left| -nK_X + lF \right|$, for some $n > 0$, such that the pair $(X, \frac{1}{n}\mathcal{M})$ is non canonical. Suppose a curve $C$ is the center of a non-canonical singularity, then $\operatorname{mult}_C \mathcal{M} = \nu > n$. Suppose $C \in F$ for some fiber $F$, then for a generic $D \in \mathcal{M}$,

$$n = D|_F \cdot (-K_F) = \nu C \cdot (-K_F) + D' \cdot (-K_F) \geqslant \nu > n.$$

Assume that $C$ is not in a fiber, then for a generic fiber $F$ let $P \in F \cap C$. Since $F$ is generic and $\mathcal{M}$ is mobile, the system $\mathcal{M}|_F$ does not have fixed components. The point $P$ is a base point of multiplicity $\nu > n$ for $\mathcal{M}|_F \in \left| -nK_F \right|$, but for generic curves $Z_1, Z_2 \in \mathcal{M}|_F$

$$n^2 = Z_1 \cdot Z_2 \geqslant (Z_1 \cdot Z_2)_x \geqslant \nu^2.$$

Now suppose that a nonsingular point $P \in X$ is the center on non-canonical singularity of $(X, \frac{1}{n}\mathcal{M})$ and let $F$ be a fiber containing $P$. Assume $P$ is an ordinary double point of $F$ and consider the blow up $f : X^+ \to X$ of $P$. For a generic $D \in \mathcal{M}$ we have

$$K_{X^+} + \frac{1}{n} D^+ (a-2)E = f^*(K_X + \frac{1}{n} D),$$

where $D^+$ is a proper transform of $D$ and $E$ is the exceptional divisor. Note that $a > 1$ since the pair $(X, \frac{1}{n}D)$ is not canonical at $P$. Consider a generic curve $C \in \left| -2K_F \right|$ passing through $P$. Because $P$ is an ordinary double point of $F$, the intersection of proper transform $C^+$ of $C$ with $E$ equals 2. Since $f^*(D) = D^+ + anE$ we get the following

$$0 \leqslant D^+ \cdot C^+ = f^*(D) \cdot C^+ - anC^+ \cdot E = D \cdot C - 2na = 2n(1-a) < 0,$$

contradiction.

If $P$ is not an ordinary double point of a fiber $F$, then for a generic $D \in \mathcal{M}$ the point $P$ is a non-log canonical center of the pair $(F, \frac{1}{n}D)$. It follows from the Corollary 4.6 that $P$ must be the cusp of some curve of anticanonical degree 1, but it is impossible by construction.

Let $P$ be an ordinary double point of $X$ and suppose that $P$ is the center on non-canonical singularity of $(X, \frac{1}{n}\mathcal{M})$. Then

$$K_{X^+} + \frac{1}{n}D^+(a-1)E = f^*(K_X + \frac{1}{n}D),$$

and $a > 1$ by Lemma 4.9. Let $F$ be a fiber containing $P$ and consider a generic curve $C \in \left| -2K_F \right|$ passing through $P$. Since $f^*(D) = D^+ + anE$ we once again get

$$0 \leqslant D^+ \cdot C^+ = f^*(D) \cdot C^+ - anC^+ \cdot E \leqslant D \cdot C - 2na = 2n(1-a) < 0.$$

□

# 5 Rationally connected non-Fano type varieties

**Proposition 5.1.** *Let $X$ be a variety of Fano type. Then there is a Mori fiber space $V$ birationally equivalent to $X$ such that $-K_V$ is big.*

*Proof.* There is an effective divisor $D$ on $X$ such that $-(K_X + D)$ is ample and $(X, D)$ is klt. Since $-K_X$ is a sum of an effective and an ample divisor it is big. If $X$ is not $\mathbb{Q}$-factorial we may replace it by its $\mathbb{Q}$-factorization $Y$ ([1, Corollary 1.4.3]). Note that there is a morphism $\pi : Y \to X$ which is an isomorphism in codimension 1, therefore $-K_Y$ is big and $(Y, D_Y)$ is klt. If $X$ is $\mathbb{Q}$-factorial, set $Y = X$. Since $-K_Y$ is $\mathbb{Q}$-Cartier and $(Y, D_Y)$ is klt, $Y$ is log terminal. There is morphism $f : Z \to Y$ such that $Z$ is terminal and all exceptional divisors $E_i$ of $f$ have $a(E, X) \leqslant 0$ ([1, Corollary 1.4.3]). The anticanonical class of $Z$ is big since

$$-K_Z = -f^*K_Y - \sum a(E_i, Y)E_i.$$

Suppose $V$ is a result of running the MMP on $Z$. For a divisorial contraction $h : W \to U$ with the exceptional divisor $E$ we can write

$$-h^*K_U = -K_W + aE, \quad a > 0,$$

hence $-K_V$ is big. The variety $X$ is rationally connected by [23] therefore $V$ is a Mori fiber space. □

*Proof of Corollary 1.2.* It follows from Theorem 1.1 and Lemma 3.8 and Lemma 4.6 that there are no birational maps from $V$ and $X$ to a Mori fiber space with big anticanonical class. Therefore, by Proposition 5.1, $V$ and $X$ are not birationally equivalent to a variety of Fano type. □

**Theorem 5.2** ([12, Theorem 0.1])**.** *Any smooth Fano variety is rationally connected.*

**Theorem 5.3** ([7, Corollary 1.3])**.** *Let $X \to B$ be a dominant morphism. If $B$ is rationally connected and the generic fiber is rationally connected then $X$ is rationally connected.*

**Corollary 5.4.** *The varieties $V$ and $X$ described in Theorem 1.1 are rationally connected.*

*Proof.* Since $V$ and $X$ are Fano fibrations over $\mathbb{P}^1$ with smooth generic fiber, it follows from Theorems 5.2 and 5.3, that they are rationally connected. □

Thus the varieties $V$ and $X$ are the examples of rationally connected varieties which are not birational to varieties of Fano type.

**Remark 5.5.** We could also construct an example by using fibrations onto Fano hypersurfaces of index one. But it is more tiresome, a little bit harder and provides an example only for dimension $\geqslant 9$.

**Remark 5.6.** Question 1 can be asked for geometrically rational surfaces defined over non-closed fields, the answer is not known. We can use similar ideas in dimension 2 over non-closed fields. A minimal conic bundle with negative self intersection index is birationally rigid. We can find a birationally rigid conic bundle $X$ with $-K_X$ not big, but we do not know if that property will hold under fiberwise maps. If it could be shown that there are no minimal conic bundles with big anticanonical class and negative $K^2$ then the answer to Question 1 would be negative in dimension 2 for non-closed fields.

**Remark 5.7.** We can run $D$-MMP on a Mori dream space for any divisor $D$ which is not nef [9, Proposition 1.11]. Thus, we have another class of varieties which behave very well under $D$-MMP. It has been proven in [1, Corollary 1.3.2] that every $\mathbb{Q}$-factorial variety of Fano type is a Mori dream space. The converse is not true even for smooth Mori dream spaces: there exists a smooth rational Mori dream space of dimension 2 which is not of Fano type [21, Section 3]. In fact, a $\mathbb{Q}$-factorial normal projective variety is of Fano type if and only if it is a Mori dream space and spectrum of its Cox ring has only log terminal singularities [8, Theorem 1.1]. One could ask Question 1 for Mori dream spaces instead of varieties of Fano type.

# References


[1] *C. Birkar, P. Cascini, C. D. Hacon,J. McKernan,* Existence of minimal models for varieties of log general type, *J. Amer. Math. Soc.*, 23, 405-468, 2010.

[2] *P. Cascini, Y. Gongyo,* On the anti-canonical ring and varieties of Fano type, *Saitama Math. J.*, 30, 27-38, 2013.

[3] *I. Cheltsov,* On singular cubic surfaces, *Asian J. Math*, Vol. 13, No. 2, 191-214, 2009.

[4] *D. Cox, J. Little, H. Schenck* Toric varieties, *Graduate Studies in Mathematics, AMS*, Vol. 124 2011.

[5] *I. Cheltsov, C. Shramov,* Log canonical thresholds of smooth Fano threefolds. With an appendix by Jean-Pierre Demailly, *Russian Mathematical Surveys*, Vol. 63:5, 859-958, 2008.

[6] *A. Corti,* Factoring birational maps of threefold after Sarkisov, *J. Algebraic geometry*, 2, 223-254, 1995.

[7] *T. Graber, J. Harris, J. Starr.* Families of rationally connected varieties, *J. Amer. Math. Soc.*, 16, 57-67, 2003.

[8] *Y. Gongyo, S. Okawa, A. Sannai, S. Takagi,* Characterization of varieties of Fano type via singularities of Cox rings, *J. Algebraic Geom*, 24, 159-182, 2015.

[9] *Y. Hu, S. Keel,* Mori dream spaces and GIT, *Michigan Math. J.*, 48, 331-348, 2000.

[10] *J. Park,* Birational maps of del Pezzo fibrations, *Crelles Journal*, vol. 2001, 538, 213-221, 2006.

[11] *J. Kollar,* Flips and abundance for algebraic threefolds, *Asterisque*, 211, Soc. Math. France, Paris, 1992.

[12] *J. Kollar, Y. Miyaoka, S. Mori,* Rationally connected varieties, *J. Alg. Geom.*, 1, 429-448, 1992.

[13] *J. Kollar,* Singularities of pairs, *E-Print*, arXiv:alg-geom/9601026v2, 1995.

[14] *Yu. G. Prokhorov,* Lectures on complements on log surfaces, *MSJ Memoirs*, 10, Math. Soc. of Japan, Tokyo, 2001.

[15] *Yu. G. Prokhorov, V. V. Shokurov,* Towards the second main theorem on complements, *J. Algebraic Geom.*, 18, no. 1, 151-199, 2009.

[16] *A. V. Pukhlikov,* Essentials of the method of maximal singularities, *London Math. Soc. Lec. Notes Series*, 281, 73-100, 2000.

[17] *A. V. Pukhlikov,* Birational geometry of Fano direct products, *Izvestiya: Math.*, 69, No. 6, 1225-1255, 2005.

[18] *A. Pukhlikov,* Birational geometry of singular Fano varieties, *Proc. Steklov Inst. Math.*, 264, No. 1, 159-177, 2009.

[19] *A. V. Pukhlikov,* Birationally rigid varieties. II. Fano fiber spaces, *Russian Math. surveys*, 65, No. 6, 1083-1171, 2010.

[20] *S. Rams,* Defect and hodge numbers of hypersurfaces, *Adv. Geom.*, 8, 257-288, 2008.

[21] *D. Testa, A. Varilly-Alvarado, M. Velasco,* Big rational surfaces, *Math. Ann.*, 351, 95-107, 2011.

[22] *G. Tian,* On Kähler–Einstein metrics on certain Kähler manifolds with $c_1(M) > 0$, *Inventiones Mathematicae*, 89, 225-246, 1987.

[23] *Q. Zhang,* Rational connectedness of log $\mathbb{Q}$-Fano varieties, *J. Reine Angew. Math.*, 590, 131-142, 2006.